

Characteristics of the UNC Kidney Center Screening Population as of 06/12/2008 (N=1,530)

Characteristics	Overall (N=1530)	Normoalbuminuria (N=548)	Microalbuminuria** (N=745)	P-value***
Age, mean (STDV)	54 (16.3)	54 (16.0)	53 (16.5)	0.201
Female, %	70	73.6	65.5	0.002
Race/Ethnicity, %				0.037
White	37	40.1	32.2	
Black	47	41.8	50.0	.
Hispanic	14	16.5	15.4	.
Native American	<1	0.2	0.4	.
Asian	1	0.5	0.8	.
Other	1	0.9	1.2	.
High School of More, %	79	75.6	79.1	0.155
Smoke 100 cigs in life, %	65	64.6	61.7	.
BMI, mean, (STDV)	30 (15.1)	29 (6.7)	31 (19.0)	0.009
Obese*%	42	36.1	46.0	<.001
Has Health Insurance, %	84	84.4	81.8	0.229
Diabetes, %	22	18.8	25.4	0.005
Hypertension,%	46	43.9	49.0	0.078
Family History of Hypertension, %	78	75.2	78.7	0.185
Family History of Diabetes, %	66	66.1	67.4	0.651
Family History of Kidney Disease,%	23	27.6	23.2	0.105
NC County, %				.
Anson	10	8.2	9.5	
Bertie	8	10.9	7.8	.
Davidson	2	0.9	2.7	.
Edgecombe	6	2.4	6.4	.
Greene	6	6.4	3.5	.
Hertford	3	1.6	4.3	.
Martin	8	8.4	9.3	.
Montgomery	33	32.7	28.6	.
Mecklenburg	7	6.2	8.1	.
Orange	15	19.0	15.8	.
Richmond	1	1.6	1.3	.
Stanley	2	1.6	2.7	.
Microalbuminuria***%	58			.

* Obesity defined as BMI>30kg/m².

**Microalbuminuria defined as urine albumin concentration>20mg/L (using semi-quantitative Micral strips).
There are 237 observations missing Microalbuminuria.

*** P-values reported are from Fishers Exact test for all categorical variables and from t-tests for all continuous variables.

Items on long-survey only (N=1358)